

### 金森先生によるセミナー（10月5日 16：20～）


[ [Topics](#) ] 2017年10月02日

リーディング大学院生の皆様  
関係教員の皆様

金森 博雄 先生（カリフォルニア工科大学）によるリーディング大学院生向けのセミナーを、以下の通り開催します。

2011年東北地方太平洋沖地震のような“Outlier（異常値）”とも言える巨大地震・津波がなぜ発生するのか、最新の知見を交えてお話いただきます。

皆様ぜひご参加ください。

[金森先生によるセミナー](#)（10月 5日 16：20 ～ 17：20） 

G-Safety Seminar: “Learning from Outliers”

Prof. Hiroo Kanamori (Seismological Laboratory, California Institute of Technology)

October 5 (Thu), 2017, 16:20 ~ 17:20

Leading Lecture Room, 9th Floor, Engineering Laboratory Complex Building, Graduate School of Engineering ([Building “C-10” in the map](#))

# “Learning from Outliers”

**Prof. Hiroo Kanamori**

(Seismological Laboratory, California Institute of Technology)

October 5 (Thu), 2017, 16:20~17:20

Leading Lecture Room, 9th Floor, Engineering Laboratory  
Complex Building, Graduate School of Engineering

Seismologists study every earthquake and try to extract source parameters. These studies form the basis of kinematic seismology, and provide key information on the general characteristics of earthquakes. The results are often presented as scaling relations between different source parameters. However, these scaling relations give only the general, or average, characteristics of earthquakes, and we often experience significant outliers. Examples are the 2004 Sumatra earthquake ( $M_w=9.1$ ), the 1992 Nicaragua tsunami earthquake ( $M_w=7.7$ ), and the 2011 Tohoku earthquake ( $M_w=9.0$ ). These outliers are often most damaging and we need to understand why they do not follow the general rule. The 2004 Sumatra and the 1992 Nicaragua earthquakes ruptured over a large distance by successively triggering many asperities. Triggering is a stochastic process that is controlled by many factors and can be predicted only statistically. Thus, a simple general rule does not work. The 2011 Tohoku earthquake had unusually large slip and stress drop near the trench, and our knowledge about the properties of sediments is still very limited. Other examples of outliers may be the large long-period ground motions that occurred in the Osaka basin during the 2011 Tohoku earthquake and in Kathmandu during the 2015 Nepal earthquake. For comprehensive understanding of seismic hazard, it is important to understand why outliers occur. We will summarize outliers and discuss why they occur in light of recent studies.